



## 저작자표시-비영리-변경금지 2.0 대한민국

이용자는 아래의 조건을 따르는 경우에 한하여 자유롭게

- 이 저작물을 복제, 배포, 전송, 전시, 공연 및 방송할 수 있습니다.

다음과 같은 조건을 따라야 합니다:



저작자표시. 귀하는 원저작자를 표시하여야 합니다.



비영리. 귀하는 이 저작물을 영리 목적으로 이용할 수 없습니다.



변경금지. 귀하는 이 저작물을 개작, 변형 또는 가공할 수 없습니다.

- 귀하는, 이 저작물의 재이용이나 배포의 경우, 이 저작물에 적용된 이용허락조건을 명확하게 나타내어야 합니다.
- 저작권자로부터 별도의 허가를 받으면 이러한 조건들은 적용되지 않습니다.

저작권법에 따른 이용자의 권리는 위의 내용에 의하여 영향을 받지 않습니다.

이것은 [이용허락규약\(Legal Code\)](#)을 이해하기 쉽게 요약한 것입니다.

[Disclaimer](#)

이학석사 학위논문

**Investigation of co-occurrence patterns of wood-  
decaying fungi and ants in dead pines of South  
Korea**

**한국의 죽은 소나무에서 공동 발생하는 목재부후균과  
개미에 대한 고찰**

2019 년 6 월

서울대학교 대학원

생명과학부

**Lupala, Abel**

# **Investigation of co-occurrence patterns of wood-decaying fungi and ants in dead pines of South Korea**

**Advisor: Professor Young Woon Lim, Ph.D.**

**A thesis submitted in partial satisfaction of the  
Requirements for the degree Master of Sciences in  
Biological sciences**

**June 2019**

**Graduate School of Biological Sciences**

**Seoul National University**

# **Investigation of co-occurrence patterns of wood-decaying fungi and ants in dead pines of South Korea**

**Lupala Abel**

**School of Biological Sciences**

**The graduate school**

**Seoul National University**

## **Abstract**

Interaction between fungi and insects such as ants, beetles, wasps and termites inhabiting dead pines has significant ecological implication in the forest as they can decompose wood debris and add nutrients to the soil; however, only scarce information is available regarding the interaction between wood-decaying fungi and ants. Wood-decaying fungi co-occurring with ants in dead pines of South Korea were investigated across 11 localities. A total of 57 pairs of wood-decaying fungi and ants were collected. 30 species of wood-decaying fungi and 14 species of ants were identified based on morphology and molecular analysis. Fungal species belonging to *Trichaptum*, *Xylodon*, *Hyphodontia*, and *Ceriporia* were dominant and co-occurred with common ant species of *Lasius*, *Camponotus*,

*Pristomyrmex*, and *Crematogaster* across the multiple sampling sites. This study provides a new baseline in unravelling the complex interaction between wood-decaying fungi and ants in forest ecosystems.

**Keywords:** Ants, Co-occurrence, Decayed pinewood, *Pinus densiflora*, Wood-decaying fungi

**Student number:** 2017-27643

## List of contents

<b>Abstract</b> .....	i
<b>List of contents</b> .....	iii
<b>List of Figures</b> .....	iv
<b>List of Tables</b> .....	v
<b>1. Introduction</b> .....	1
1.1. Wood-decaying fungi.....	1
1.2. Wood-decaying fungi and insects' interactions .....	1
1.3. Korean pine trees .....	3
1.4. Objective .....	4
<b>2. Materials and methods</b> .....	5
2.1. Sampling sites and specimen collection .....	5
2.1. Fungi collection and identification .....	8
2.3. Ants collection and identification .....	9
<b>3. Results</b> .....	13
3.1. Wood-decaying fungi inhabiting dead pinewoods .....	13
3.2. Ants nesting in dead pinewoods .....	21
3.3. Co-occurrence patterns of wood-decaying fungi and ants.....	28
<b>4. Discussion</b> .....	31
<b>5. Conclusions</b> .....	35
<b>6. References</b> .....	36
<b>7. Abstract in Korean</b> .....	45

## List of Figures

<b>Figure 1.</b> Map of South Korea showing sampling localities.....	6
<b>Figure 2.</b> Sampling approach.....	7
<b>Figure 3.</b> Location of Primers for PCR amplification. A- ITS region, B- COI region.....	12
<b>Figure 4.</b> Representative list of collected wood-decaying fungi. A-L.....	19
<b>Figure 5.</b> Pie charts of collected wood-decaying fungi across the sampling sites.....	20
<b>Figure 6.</b> Collected Ants species across sampling sites. A-N.....	26
<b>Figure 7.</b> Pie charts of collected ants across the sampling sites.....	27
<b>Figure 8.</b> A flow plot of co-occurring pairs of wood-decaying fungi and ants. The left side bars represent families and species of wood-decaying fungi; the right side bars represent the ants' tribes and species. The length of a bar next to the name of a species is proportional to its number of co-occurring pairs.....	30

## **List of Tables**

<b>Table 1.</b> Information on the primers used for the PCR in this study.....	11
<b>Table 2.</b> List of collected wood-decaying fungi across sampling sites and their Genbank accession numbers.....	15
<b>Supplementary Table 1.</b> Representative list of molecular identified wood-decaying fungi species and their GenBank matches.....	17
<b>Table 3.</b> List of collected ants across sampling sites and their Genbank accession numbers.....	22
<b>Supplementary Table 2.</b> Representative list of molecular identified ant species, their Genbank and NIBR matches.....	24



# **1. Introduction**

## **1.1. Wood-decaying fungi**

Wood-decaying fungi are saprotroph and distributed across several orders of class Agaricomycetes (Basidiomycota) (Hibbett et al., 2007). These fungi can decompose wood matters into simple sugars, oligosaccharides and humic substances which are subsequently added to soil (Carlile et al., 2001). Through decomposition of fallen dead wood, these fungi provide habitats for many other organisms and regenerates forests' biodiversity (Sippola et al., 2005; Heilmann-Clausen et al., 2014; Fukasawa et al., 2015). Wood decomposition is a complex ecological process (Herrmann and Bauhus, 2013) which is regulated by different factors: abiotic (e.g. climate conditions) and biotic (e.g. substrate quality) factors and the composition and activity of decomposer fungal communities (Weedon et al., 2009; Liu et al., 2013).

## **1.2. Wood-decaying fungi and insects' interactions**

Habitats and nutrients generated by wood-decaying fungi are sufficient factors to attract various invertebrates (Johansson et al., 2006; Warren and Bradford, 2011). Some of them have also developed symbiotic relationships with these fungi (Hågvar, 1999; Mueller et al., 2005; Schigel, 2011a; Li et al., 2015). Wood boring ambrosia beetles (*Ambrosiodmus* and *Ambrosiophilus*) have a mutualistic

association with the wood degrading fungus, *Flavodon ambrosius* (Hulcr and Stelinski, 2017; Li et al., 2017). Another form of association between fungi and insect is between fungus-cultivating termites (Termitidae, Macrotermitinae) and their obligate mutualistic fungi (Agaricomycetes, Lyophyllaceae, Termitomyces) whereby in this mutualistic association only one termite species is associated to one fungi species (Makonde et al., 2013).

Ants have also exhibited strong co-evolutionary dynamics with fungi (Masiulionis et al., 2014). The most widely known and explored ant-fungus mutualism is between the Attine ants (tribe Attini), best known as “leaf cutter ants” (*Atta* and *Acromyrmex* genera) and their mutualistic fungal cultivars, Leucocoprineae and Pterulaceae, in Southern and Central America and southern parts of USA (Hinkle et al., 1994; Mehdiabadi and Schultz, 2010; Masiulionis et al., 2014). Masiulionis et al (2014) reported that ants can cultivate the fungi in the nest and protect the fungi from alien microorganism such as bacteria and other invading fungi and, in return, these fungi provide nutrients to the ants in the form of specialized inflated hyphal tips called gongylidia. Ants of diverse genera have been observed as common visitors of wood-decaying fungi as well (Lewis and Worthen, 1992; Orr and Charles, 1994; Mueller et al., 2001). Recently, Epps and Penick (2018) scrutinized mushroom visiting ants (*Aphaenogaster* spp.) and reported that these ants with a generalist diet fed on wood-decaying fungus, *Pleurotus ostreatus*.

### 1.3. Korean pine trees

Coniferous forests have been dominating since 1970 (Kwon., 2016). Out of major conifers such as *Pinus densiflora* (Korean red pine), *Pinus koraiensis* (Korean pine), *Pinus thunbergii* (Black pine), and *Pinus rigida* (Pitch pine), *Pinus densiflora* is the most abundant one in South Korea. This tree is native to Korea, Japan, China, and East Russia (Yu et al., 2004). According to the Korea Forest Service 2016; *P. densiflora* accounts for the 87% of the species in coniferous forests in the country. It has an immense ecological as well as economic importance since it is used for the production of timber as well as playing an important role in many forests' ecosystems (Kwon., 2016).

For the case of dead pines, they are nutrients and habitat source to wood-decaying fungi (Kim et al., 2017). Diverse species of wood-decaying fungi have been reported to be inhabiting dead pines, these include species like *Athelia epiphylla*, *Fomitopsis pinicola*, *Gloeophyllum sepiarium*, *Oligoporus placentus*, *Phanerochaete filamentosa*, and *Trichaptum abietinum* (Son et al., 2011). These wood-decaying fungi attract insects such as woodborers (Park et al., 2014; Floren et al., 2015) and wasp (Pažoutová and Šrůtka, 2007).


#### **1.4. Objective**

Several species of wood-decaying fungi have mutualistic interactions with beetles and wasps (Son et al., 2011; Park et al., 2014., Rajala et al., 2015). In South Korea, where numerous species of wood-decaying fungi and ants such as *Aphaenogaster japonica*, *Camponotus japonicus*, and *Formica japonica* have long been observed to inhabit dead pinewoods (Kwon et al., 2012; Kwon et al., 2013; Kim et al., 2017). However, there has not been a comparable study of mutualistic associations between wood-decaying fungi and ants inhabiting dead pinewoods. To understand their relationship, investigation of their co-occurrence patterns is required first. Therefore, the main objective of this study is to investigate the co-occurrence patterns of wood-decaying fungi and ants in dead pinewoods. Furthermore, to assess which ant-fungi pair have the highest co-occurrence.

## **2. Materials and methods**

### **2.1. Sampling sites and specimen collection**

Sampling of wood-decaying fungi and co-occurring ants on dead pinewoods was conducted between mid-March to September 2018 across 57 sites of 11 locations in Korea (Fig. 1). They were collected only after confirming that (1) wood-decaying fungi and ants were both present on a dead pinewood and (2) an ant nest was found within the dead pinewood. Based on these criteria, samples were not collected when ants simply foraged wood-decaying fungi without a nest underneath. Fungi and ants were collected in pairs after they were confirmed to co-exist in a dead pinewood, either standing or fallen (Fig. 2). All sites were of lower to higher altitude mountainous regions with moderate understory vegetation to completely covered litter layers. In all sites, vegetation was dominated by *Pinus densiflora*.



**Figure 1.** Map of South Korea showing sampling localities.

## Ant nest and fungi



**Wood-decaying fungi**



**Ant**

**Figure 2.** Sampling approach. Collection of co-existing wood –decaying fungi and ants from a dead pine pinewood with ants' nest.

## **2.1. Fungi collection and identification**

Fruiting bodies were collected, dried and deposited in the Seoul National University Fungus Collection (SFC). They were identified via morphological and molecular approaches. Classical methodology was achieved by macro- and micro-morphological observations using taxonomic guides (Largent and Thiers, 1977; Breitenbach and Kränzlin, 1986)

Genomic DNA was extracted from fruiting bodies using a modified Cetyl trimethyl ammonium bromide (CTAB) extraction protocol of Rogers and Bendich (1994). The internal transcribed spacer (ITS) (Fig. 3A) region was amplified using combinations of the forward primers (ITS1F and NS11) and reverse primers (ITS4B, LB-W, and NLB4) (Table 1) (White et al., 1990; Gardes and Bruns, 1994; Martin and Rygiewicz, 2005; Tedersoo et al., 2008). PCR amplification was performed as described by Park et al. (2013). PCR products were visualized on a 1% agarose gel and purified using the Expin PCR purification kit (GeneAll Biotechnology, Seoul, South Korea). Sanger sequencing was performed at Macrogen (Seoul, South Korea) on an automated DNA sequencer (ABI Prism 3730XL analyzer; Applied Biosystems, Foster City, California) using the aforementioned PCR primers.



Newly generated sequences were proofread and edited using MEGA 5 (Tamura et al., 2011). Molecular identification was conducted with ITS region using BLAST against GenBank database. Species level identification was achieved through the sequence similarity with 98 to 100%. Specimens that had sequence similarity of 95 to 97% or no species information in open database remained at the genus level (Table S1). All sequences were deposited at GenBank under accession number MK992816- MK992868 (Table 2).

### **2.3. Ants collection and identification**

Ants were collected into 1.5 mL Eppendorf tubes containing 70% ethanol by employing hand collection method and transported to the laboratory. Their identification was accomplished by morphological and molecular approaches. For morphological approach, all collected ants were observed under Nikon SMZ1500 stereo microscope and identified according to the taxonomic keys (Japanese Ant Image Database, 2010 (<http://ant.edb.miyakyo-u.ac.jp/>); Kwon et al., 2012; Dong, 2017).

Molecular approach was performed as follows. An individual ant was placed in a 1.5 mL Eppendorf tube after the removal of gaster which is known to contain a substance inhibiting the PCR (Feldhaar et al., 2003). Genomic DNA was

extracted by using AccuPrep® Genomic DNA Extraction Kit (Bioneer, Daejeon, South Korea) according to the manufacturer's protocol. For PCR amplification and sequencing, the insect barcode mitochondrial cytochrome c oxidase subunit I (COI) gene was targeted (Fig. 3B) which is the basis of ants' molecular identification (Wilson, 2012). COI was amplified using the primers LCO1490 and HCO2198 (Table 1) (Folmer et al., 1994). PCRs were conducted in the C1000 thermal cycler (Bio-Rad, Hercules, CA, USA) using the AccuPower PCR Premix (Bioneer Co., Daejeon, Korea) in a final volume of 20 µL containing 10 pmol of each primer and 1 µL of DNA under the following conditions: 95°C for 5 min, followed by 35 cycles of 95°C for 40 secs, 45°C for 40 secs, and 72°C for 1 min and a final extension step at 72°C for 10 min.

Purification and sequencing of PCR product was conducted as described in the previous section. As reference sequences of ants native to East Asia are largely unavailable in GenBank, for some ants' specimens, identification was achieved through the BLAST service of the National Institute of Biological Resources (<https://species.nibr.go.kr>) (Table S2). All ants' sequences were deposited at GenBank under accession number MN010595-MN010641 (Table 3).

**Table 1. Information on the primers used for the PCR in this study**

<b>Region</b>	<b>Primer</b>	<b>Sequence (5'-3')</b>	<b>Reference</b>
<b>For Fungi</b>			
<b>ITS</b>	NSI1	GATTGAATGGCTTAGTGAGG	Martin and Rygiewicz, 2005
	NLB4	GGATTCTCACCTCTATGAC	Martin and Rygiewicz, 2005
	ITS1F	CTTGGTCATTTAGAGGAAGTAA	White et al., 1990
	ITS4B	CAGGAGACTTGTACACGGTCC	Gardes and Bruns, 1994
	LB-W	CTTTTCATCTTTCCTCACGG	Tedersoo et al., 2008
<b>For Ants</b>			
<b>COI</b>	LCO1490	GGTCAACAAATCATAAAGATATTGG	Folmer et al., 1994
	HCO2198	TAAACTTCAGGGTGACCAAAAAATCA	Folmer et al., 1994



### 3. Results

#### 3.1. Wood-decaying fungi inhabiting dead pinewoods

57 wood-decaying fungi co-inhabited with ants were collected from dead pinewoods. They were identified by complementing both morphological and molecular analyses. Five fungal species were first identified morphologically based on their distinctive characteristics followed by molecular confirmation: *Ceriporia nanlingensis*, *Heterobasidion ecrustosum*, *Schizophyllum commune*, *Trichaptum abietinum*, and *Xylodon flaviporus* (Fig. 4). Other species were identified first based on ITS sequence analysis and then confirmed by morphological characters. Forty-eight samples were identified to species level and seven to genus level while two to order level only. A total of 30 species of wood-decaying fungi was confirmed from 57 sites (Table 2).

At the family level, species of Polyporaceae and Schizoporaceae were the most dominant, followed by Meruliaceae, Rickenellaceae, Tubulicrinaceae, Botryobasidiaceae, and Fomitopsidaceae. The remaining seven families were mostly collected from single sites only (Fig. 5). At the genus level, *Xylodon* and *Trichaptum* were the most dominant genera followed by *Hyphodontia*, *Ceriporia*, and *Botryobasidium* (Fig. 5). In terms of species, *T. abietinum* had the highest collection frequency (9 sites) followed by *Xylodon ovisporus*, *Resinicium*

*furfuraceum*, and *Hyphodontia pallidula* (each 5 sites). *Ceriporia alachuana* and *Ceriporia naglingensis* were collected from five sites. Species such as *Amylostereum areolatum*, *Heterobasidion ecrustosum*, *Perenniporia koreana*, *Phlebia acanthocystis*, *Postia hirsuta* and *Pseudomerulius aureus* were collected only once (Table 2).

**Table 2: List of collected wood-decaying fungi across sampling sites and their Genbank accession numbers**

Wood-decaying fungi	Specimen code	Locality	ITS Accession No.
<b>Amylostereaceae</b>			
<i>Amylostereum areolatum</i>	SFC20180314-02	1	MK992817
<b>Atheliaceae</b>			
<i>Athelia epiphylla</i>	SFC20180314-01	1	MK992816
<b>Bondarzewiaceae</b>			
<i>Heterobasidion ecrustosum</i>	SFC20180919-03	11	MK992864
<b>Botryobasidiaceae</b>			
<i>Botryobasidium</i> sp. 1	SFC20180710-21	7	MK992837
<i>Botryobasidium</i> sp. 2	SFC20180609-01	4	MK992824
	SFC20180710-16	6	MK992832
<i>Botryobasidium</i> sp. 3	SFC20180710-22	7	MK992838
<b>Fomitopsidaceae</b>			
<i>Dacryobolus</i> sp. 1	SFC20180907-154	9	MK992862
<i>Dacryobolus</i> sp. 2	SFC20180907-153	9	MK992861
<i>Postia hirsuta</i>	SFC20180822-19	4	MK992857
<b>Hydnodontaceae</b>			
<i>Trechispora confinis</i>	SFC20180710-18	6	MK992834
	SFC20180710-23	7	MK992839
<b>Meruliaceae</b>			
<i>Ceriporia alachuana</i>	SFC20180710-15	6	MK992831
	SFC20180810-01	4	MK992849
	SFC20180810-03	4	MK992851
<i>C. nanlingensis</i>	SFC20180710-17	6	MK992833
	SFC20180822-20	4	MK992858
<i>Phlebia acanthocystis</i>	SFC20180704-91	5	MK992826
<i>P. acerina</i>	SFC20180711-01	8	MK992841
<b>Phanerochaetaceae</b>			
<i>Phlebiopsis gigantea</i>	SFC20180711-02	8	MK992842
<b>Polyporaceae</b>			
<i>Perenniporia koreana</i>	SFC20180719-04	4	MK992847
<i>Trichaptum abietinum</i>	SFC20180531-03	2	MK992820
	SFC20180531-04	2	-
	SFC20180710-12	6	MK992828
	SFC20180710-13	6	MK992829
	SFC20180710-14	6	MK992830
	SFC20180719-05	4	MK992848
	SFC20180822-21	4	-
	SFC20180919-01	11	-

**Table 2** Continued

	SFC20180928-35	1	-
<b>Rickenellaceae</b>			
<i>Peniophorella pubera</i>	SFC20180601-02	3	MK992822
<i>P. subpraetermissa</i>	SFC20180719-01	4	MK992844
<i>Resinicium furfuraceum</i>	SFC20180822-18	4	MK992856
	SFC20180907-152	9	MK992860
	SFC20180919-02	11	MK992863
	SFC20180919-06	11	MK992867
	SFC20180929-32	1	MK992868
<b>Schizophyllaceae</b>			
<i>Schizophyllum commune</i>	SFC20180704-92	5	MK992827
<b>Schizoporaceae</b>			
<i>Xylodon chinensis</i>	SFC20180710-20	6	MK992836
	SFC20180818-36	4	MK992854
<i>X. flaviporus</i>	SFC20180710-24	7	MK992840
<i>X. nespori</i>	SFC20180601-03	3	MK992818
	SFC20180822-17	4	MK992819
<i>X. ovisporus</i>	SFC20180531-01	2	MK992846
	SFC20180531-02	2	MK992850
	SFC20180719-03	4	MK992859
	SFC20180810-02	4	MK992823
	SFC20180822-22	4	MK992855
<b>Tapinellaceae</b>			
<i>Pseudomerulius aureus</i>	SFC20180818-35	4	MK992853
<b>Tubulicrinaceae</b>			
<i>Hyphodontia pallidula</i>	SFC20180601-01	3	MK992821
	SFC20180710-19	6	MK992835
	SFC20180711-03	8	MK992843
	SFC20180719-02	4	MK992845
	SFC20180810-04	4	MK992852
<i>Tubulicrinis</i> sp.	SFC20180919-05	11	MK992866
<b>Unidentified family</b>			
Auriculariales sp.	SFC20180919-04	11	MK992865
Hymenochaetales sp.	SFC20180609-02	4	MK992825




**Supplementary Table 1. Representative list of molecular identified wood-decaying fungi species and their**

**Genbank matches**


<b>Wood-decaying fungi</b>	<b>Specimen code</b>	<b>Closest fungal match (accession number)</b>	<b>Similarity (%)</b>
<i>Amylostereum areolatum</i>	SFC20180314-02	<i>Amylostereum areolatum</i> (KX449521)	100
<i>Athelia epiphylla</i>	SFC20180314-01	<i>Athelia epiphylla</i> (GU187501)	99
<i>Botryobasidium</i> sp. 1	SFC20180710-21	Cf. <i>Botryobasidium</i> (KP814186)	99
<i>Botryobasidium</i> sp. 2	SFC20180609-01	<i>Botryobasidium</i> sp. (KY462401)	99
<i>Botryobasidium</i> sp. 3	SFC20180710-22	<i>Botryobasidium</i> sp (KY462551)	100
<i>Ceriporia alachuana</i>	SFC20180710-15	<i>Ceriporia alachuana</i> (JX623897)	99
<i>Ceriporia nanlingensis</i>	SFC20180822-20	<i>Ceriporia nanlingensis</i> (JX623938)	98
<i>Dacryobolus</i> sp. 1	SFC20180907-154	<i>Dacryobolus</i> sp. (KJ668557)	99
<i>Dacryobolus</i> sp. 2	SFC20180907-153	Cf. <i>Dacryobolus</i> sp. (KP814329)	99
<i>Heterobasidion ecrustosum</i>	SFC20180919-03	<i>Heterobasidion ecrustosum</i> (KF218832)	100
<i>Hyphodontia pallidula</i>	SFC20180601-01	<i>Hyphodontia pallidula</i> (KP814392)	99
<i>Peniophorella. subpraetermissa</i>	SFC20180719-01	<i>Peniophorella. subpraetermissa</i> (NR_119600)	98
<i>Peniophorella pubera</i>	SFC20180601-02	<i>Peniophorella pubera</i> (KJ714003)	99

**Supplementary Table 1** Continued

<i>Perenniporia koreana</i>	SFC20180719-04	<i>Perenniporia koreana</i> (KJ156314)	99
<i>Phlebia acanthocystis</i>	SFC20180704-91	<i>Phlebia acanthocystis</i> (KY948767)	99
<i>Phlebia acerina</i>	SFC20180711-01	<i>Phlebia acerina</i> (KP135371)	99
<i>Phlebiopsis gigantea</i>	SFC20180711-02	<i>Phlebiopsis gigantea</i> (AF087485)	100
<i>Postia hirsute</i>	SFC20180822-19	<i>Postia hirsuta</i> (KJ684971)	99
<i>Pseudomerulius aureus</i>	SFC20180818-35	<i>Pseudomerulius aureus</i> (GU187534)	99
<i>Resinicium furfuraceum</i>	SFC20180822-18	<i>Resinicium furfuraceum</i> (KP814304)	98
<i>Schizophyllum commune</i>	SFC20180704-92	<i>Schizophyllum commune</i> (AF249378)	99
<i>Trechispora confinis</i>	SFC20180710-18	<i>Corticium confine</i> (MG231494)	98
<i>Trichaptum abietinum</i>	SFC20180710-12	<i>Trichaptum abietinum</i> (MF381023)	100
<i>Xylodon chinensis</i>	SFC20180818-36	<i>Hyphodontia chinensis</i> (KX857804)	98
<i>Xylodon ovisporus</i>	SFC20180531-02	<i>Xylodon ovisporus</i> (MF540764)	99
<i>Xylodon flaviporus</i>	SFC20180710-24	<i>Schizopora flavipora</i> (AF145585)	99
<i>Xylodon nespori</i>	SFC20180822-17	<i>Hyphodontia nespori</i> (DQ873622)	99
<i>Tubulicrinis</i> sp.	SFC20180919-05	Cf. <i>Tubulicrinis</i> sp (KP814266)	95
Auriculariales sp.	SFC20180919-04	<i>Stypella subgelatinosa</i> (MK098926)	88
Hymenochaetales sp.	SFC20180609-02	<i>Tubulicrinis globisporus</i> (DQ873655)	87



**Figure 4.** Representative list of collected wood-decaying fungi. **A-** *Ceriporia alachuana*, **B-** *Ceriporia nanlingensis*, **C-** *Dacryobolus* sp. 2, **D-** *Heterobasidion ecrustosum*, **E-** *Hyphodontia pallidula*, **F-** *Phlebia acanthocystis*, **G-** *Phlebia acerina*, **H-** *Pseudomerulius aureus*, **I-** *Resinicium furfuraceum*, **J-** *Trichaptum abietinum*, **K-** *Xylodon chinensis*, **L-** *Xylodon ovisporus*. **Scale bar = 1 cm.**



**Figure 5.** Pie charts of collected Wood-decaying fungi across the sampling sites. Inner cycle-fungal families, Outer cycle-fungal genera.

### 3.2. Ants nesting in dead pinewoods

Ants collected with wood-decaying fungi were identified as 14 different species (Fig 6) of five tribes, namely Attini, Camponotini, Crematogastrini, Lasiini (Sub-family Formicinae), and Stenammini (Sub-family Myrmicinae) (Table 3, Fig 7). Species frequently observed were *Lasius japonicus* (17 sites) and *Pristomyrmex punctatus* (9 sites), followed by *Crematogaster teranishii* (5 sites), *Pheidole fervida* (5 sites), and *Vollenhovia emeryi* (4 sites). Ants from the genus *Camponotus* had the highest species diversity: *C. atrox*, *C. kiusuensis*, *C. nipponensis*, *C. vitiosus*, and *C. quadrinotatus* (Fig 6). Species of *Aphaenogaster japonica* and *Lasius spathopus* were each collected two times while the least common species *Nylanderia flavipes* and *Polyrhachis lamellidens* were collected once (Table 3).

**Table 3: List of collected ants across sampling sites and their Genbank accession numbers**

<b>Ants</b>	<b>Specimen code</b>	<b>Locality</b>	<b>COI Accession No.</b>
<b>Attini</b>			
<i>Pheidole fervida</i>	SFC20180907-A03	9	MN010633
	SFC20180919-A03	11	MN010636
	SFC20180818-A01	4	-
	SFC20180531-A03	2	MN010598
	SFC20180531-A04	2	MN010599
<b>Camponotini</b>			
<i>Camponotus atrox</i>	SFC20180314-A01	1	MN010595
	SFC20180929-A01	1	MN010641
	SFC20180928-A01	1	MN010640
<i>C. kiusuensis</i>	SFC20180919-A01	11	MN010634
	SFC20180919-A02	11	MN010635
<i>C. nipponensis</i>	SFC20180810-A04	4	MN010624
	SFC20180822-A03	4	MN010627
	SFC20180818-A02	4	-
<i>C. quadrinotatus</i>	SFC20180822-A01	4	MN010625
<i>C. vitiosus</i>	SFC20180704-A01	5	MN010605
<i>Polyrhachis lamellidens</i>	SFC20180710-A12	7	MN010613
<b>Crematogastrini</b>			
<i>Crematogaster teranishii</i>	SFC20180710-A07	6	-
	SFC20180710-A06	6	-
	SFC20180710-A01	6	MN010606
	SFC20180710-A03	6	-
	SFC20180822-A05	4	MN010629
<i>Pristomyrmex punctatus</i>	SFC20180601-A01	3	MN010600
	SFC20180711-A01	8	MN010615
	SFC20180822-A02	4	MN010626
	SFC20180710-A02	6	MN010607
	SFC20180710-A09	6	MN010610
<i>Vollenhovia emeryi</i>	SFC20180601-A03	3	MN010602
	SFC20180531-A01	2	MN010596
	SFC20180822-A04	4	MN010628
	SFC20180609-A02	4	MN010604
	SFC20180719-A01	4	-
	SFC20180822-A06	4	MN010630

**Table 3** Continued

---

<b>Lasiini</b>			
<i>Lasius japonicus</i>	SFC20180919-A04	11	MN010637
	SFC20180710-A10	7	MN010611
	SFC20180609-A01	4	MN010603
	SFC20180710-A11	7	MN010612
	SFC20180710-A04	6	MN010608
	SFC20180810-A01	4	MN010621
	SFC20180711-A02	8	MN010616
	SFC20180719-A02	4	MN010617
	SFC20180601-A02	3	MN010601
	SFC20180719-A04	4	MN010619
	SFC20180907-A01	9	MN010631
	SFC20180919-A05	11	MN010638
	SFC20180919-A06	11	MN010639
	SFC20180710-A13	7	MN010614
	SFC20180531-A02	2	MN010597
	SFC20180719-A03	4	MN010618
<i>L. spathopus</i>	SFC20180710-A05	6	MN010609
	SFC20180810-A02	4	MN010622
<i>Nylanderia flavipes</i>	SFC20180810-A03	4	MN010623
<b>Stenammini</b>			
<i>Aphaenogaster japonica</i>	SFC20180719-A05	4	MN010620
	SFC20180907-A02	9	MN010632

---

**Supplementary Table 2. Representative list of molecular identified ant species, their Genbank and NIBR matches**

<b>Ant species</b>	<b>Code number</b>	<b>Closest match (GenBank)</b>	<b>Closest match (NIBR)</b>
<i>Aphaenogaster japonica</i>	SFC20180719-A05	<i>Aphaenogaster japonica</i> (99%) KJ920527	<i>Aphaenogaster japonica</i> (100%) WBN0014101
<i>Camponotus atrox</i>	SFC20180314-A01	<i>Camponotus atrox</i> (100%) KT159775	<i>Camponotus obscuripes</i> (100%) WBN0013599
<i>Camponotus kiusuensis</i>	SFC20180919-A01	<i>Camponotus kiusuensis</i> (99%) AB019412	<i>Camponotus kiusuensis</i> (99%) WBN0013608
<i>Camponotus nipponensis</i>	SFC20180810-A04	<i>Camponotus</i> sp (88%) KX347287	<i>Camponotus nipponensis</i> (99%) WBN0013611
<i>Camponotus quadrinotatus</i>	SFC20180822-A01	<i>Camponotus quadrinotatus</i> (98%) AB019414	<i>Camponotus quadrinotatus</i> (99%) WBN0013617
<i>Camponotus vitiosus</i>	SFC20180704-A01	<i>Camponotus vitiosus</i> (100%) AB019416	-
<i>Crematogaster teranishii</i>	SFC20180710-A01	<i>Crematogaster</i> sp (90%) KC501979	<i>Crematogaster teranishii</i> (98%) WBN0013993
<i>Lasius japonicus</i>	SFC20180710-A10	<i>Lasius japonicus</i> (98%) AB371063	<i>Lasius japonicus</i> (94.1%) WBN0013662
<i>Lasius spathepus</i>	SFC20180810-A02	<i>Lasius fuliginosus</i> (93%) AY519394	-
<i>Nylanderia flavipes</i>	SFC20180810-A02	<i>Nylanderia flavipes</i> (99%) MH754327	-
<i>Pheidole fervida</i>	SFC20180907-A03	<i>Pheidole fervida</i> (98%) MH754425	<i>Pheidole fervida</i> (99.5%) WBN0014074
<i>Polyrhachis lamellidens</i>	SFC20180710-A12	<i>Polyrhachis lamellidens</i> (99%) KM348369	<i>Polyrhachis lamellidens</i> (100%) WBN0013700




**Supplementary Table 2** Continued

<i>Pristomyrmex punctatus</i>	SFC20180601-A01	<i>Pristomyrmex punctatus</i> (99%) AB556946	-
<i>Vollenhovia emeryi</i>	SFC20180822-A04	<i>Vollenhovia emeryi</i> (99%) KF815843	<i>Vollenhovia emeryi</i> (98%) WBN0013981
-No sequence information			



**Figure 6.** Collected Ants species across sampling sites. **A-** *Aphaenogaster japonica*, **B-** *Camponotus atrox*, **C-** *Camponotus kiusuensis*, **D-** *Camponotus nipponensis*, **E-** *Camponotus quadrinotatus*, **F-** *Camponotus vitiosus* **G-** *Crematogaster teranishii*, **H-** *Lasius japonicus*, **I-** *Lasius spathepus*, **J-** *Nylanderia flavipes*, **K-** *Pheidole fervida*, **L-** *Polyrhachis lamellidens*, **M-** *Pristomyrmex punctatus*, **N-** *Vollenhovia emeryi*. Scale bar = 1 mm.



**Figure 7.** Pie charts of collected ants across the sampling sites. Inner circle ants' tribes, Outer circle ants' genera.


### 3.3. Co-occurrence patterns of wood-decaying fungi and ants

A total of 57 pairs of wood-decaying fungi and ants were detected from 11 locations. Ants' nests were deep inside dead pinewoods or near the top layer of the dead pine harboring fruiting bodies or mycelia of fungi. At the family level for fungi and tribal level for ants, Polyporaceae co-occurred with all ants' tribes. Meruliaceae, Rickenellaceae, Schizoporaceae and Tubulicrinaceae co-occurred with three tribes Camponotini, Crematogastrini and Lasinii while all Botryobasidiaceae co-occurred with tribe Lasinii only (Fig. 8).

*Trichaptum abietinum* had the highest number of co-occurrence pairs with numerous species of ants across the sampling sites: three pairs with *Crematogaster teranishii*, two pairs with *Pheidole fervida*, two pairs with *Camponotus* spp. and one pair each with *Aphaenogaster japonica* and *Pristomyrmex punctatus* (Fig. 8). Fungal species of *Xylodon*, *Botryobasidium* spp, *Ceriporia*, *Hyphodontia pallidula*, and *Resinicium furfuraceum* followed in terms of co-occurrences pairs with multiple ant species, they showed co-occurrence with *Lasius japonicus* ants. *L. japonicus* ants had the highest number of co-occurrence pairs with wood-decaying fungi across the sampling sites followed by *Pristomyrmex punctatus*, they also co-occurred with multiple fungi species across the sites, the most being three pairs with *Xylodon* species.

Co-occurrence of some wood-decaying fungi and ants was observed by the tendency of ants displaying foraging-like movements on fruiting bodies. Ants from inside the nest and outside surroundings were observed to be displaying these movements on the surface of fruiting bodies. Pairs of wood-decaying fungi and ants involved in this tendency were *Ceriporia alachuana* / *Lasius japonicas* (Fig.2), *Heterobasidion ecrustosum* / *Pheidole fervida*, *T. abietinum* / *C. teranishii*, and *X. ovisporus* / *P. punctatus*. Most of the remaining wood-decaying fungi were collected once or twice and co-occurred with commonly observed ants' species across the sites (Fig.8).

## Wood-decaying fungi



**Figure 8.** A flow plot of co-occurring pairs of wood-decaying fungi and ants. The left side bars represent families and species of wood-decaying fungi; the right side bars represent the ants' tribes and species. The length of a bar next to the name of a species is proportional to its number of co-occurring pairs.

## 4. Discussion

Wood-decaying fungi co-occurring with ants in dead pinewoods were surveyed across 11 locations in South Korea. This is the first descriptive study addressing multiple species co-occurrences (wood-decaying fungi and ants) in dead pinewoods of South Korea. Outside South Korea, many studies regarding wood-decaying fungi interactions with insects have only focused on various groups of fungivorous insects; the Czech Republic (Kula et al., 1999), the Fennoscandian countries (Hågvar, 1999; Thunes et al., 2000; Jonsell and Nordlander, 2004; Schigel, 2011b), China (Komonen et al., 2003), Russia (Nikitsky and Schigel, 2004; Krasutskii, 2007), New Zealand (Osawa et al., 2011), Borneo (Yamashita et al., 2015), and Brazil (Valer et al., 2016).

Wood-decaying fungi are key players in forest ecosystems because of their ability to decompose dead wood, recycling nutrients and initiating a successional dynamic for saproxylic arthropods (Jönsson et al., 2008; Lonsdale et al., 2008). A list of collected wood-decaying fungi from decaying pinewoods in this study was consistent with the relevant previous studies in Korea (Kim et al., 2009; Jang et al., 2014; Jang et al., 2016; Kim et al., 2017) and also outside Korea (Kubartová et al., 2012; Walker et al., 2014; Ottosson et al., 2015; Fukasawa and Matsuoka, 2015). *Athelia epiphylla* and *Schizophyllum commune* are known to be exclusive to deciduous trees (Allen et al 2000; Takemoto et al., 2010), but rarely reported from

coniferous trees such as pine (Kim et al., 2009). *Trichaptum abietinum*, which was frequently observed on pinewoods, is known as the first colonizer on dead pinewoods in many forests (Weslien et al., 2011). *Hyphodontia* and *Xylodon* (specifically *Xylodon ovisporus*) were also dominant on decaying pinewoods across the sampling sites and have been reported to be common in deciduous forests, as well (Gafforov et al., 2017). *Ceriporia* species which can cause decaying of both deciduous and coniferous wood were also common (Wulandari et al., 2018). *Ceriporia alachuana* and *Ceriporia nanlingensis* are commonly isolated from pinewood (Dai, 2012). Other observed species were all frequently found from decaying pinewood (Lee and Jung, 2006; Wooding et al., 2013; Jang et al 2016).

Ants associated with the occurrence of wood-decaying fungi in this study were 14 species. These species were all reported in previous ants' communities' surveys in South Korea (Kim and Lyu, 2012; Kwon et al., 2013, 2014a, 2014b; Kwon, 2015a; Kwon, 2015b; Kwon and Lee, 2015; Kwon, 2016). *Lasius japonicus* was the most frequently detected ant species in this study. It is frequently found in Korean forests and dead pinewood is one of its preferred nesting place (Kwon and Lee, 2015). The least collected ant species in this study, *Nylanderia flavipes* also has been reported as one of the most common ant species in Korean pine forests (Kwon and Lee, 2015). Remaining species such as *Aphaenogaster japonica*, *Camponotus* spp., *Crematogaster teranishi*, and *Pheidole fervida* have been



previously observed to be nesting inside decayed pinewood (Kwon et al., 2014b; Kwon, 2016).

For some wood-decaying fungi, interaction with insects is crucial for their establishment and colonization in dead woods (Strid et al., 2014). Insects such as bark beetles and wasps are one of the key players in dispersal of fungal spores to new dead trees (Floren et al., 2015). Bark beetles are usually the first species to colonize newly dead or dying tree and feed on the nutrient-rich phloem (Ejrnæs et al., 2006; Son et al., 2011; Weslien et al., 2011), followed by wood-decaying fungi that colonize broken or cut stems or scars in the bark (Son et al., 2011). Most of the wood-decaying fungi found in this study were previously reported to be associated with beetles and wasps (Ševčík, 2003; Persson et al., 2011; Son et al., 2011; Olatinwo et al., 2013; Strid et al., 2014). *Amylostereum areolatum* and *Athelia epiphylla* are well known decaying fungi to have an interaction with wasp and beetles, respectively (Rajala et al., 2015; Seibold et al., 2019).

Although with limited sampling, results from this study showed that various wood-decaying fungi were co-occurring with ants on dead pines. Ants were also observed to display foraging like movement on basidiocarps.

Foraging activities of *Camponotus* species on basidiocarp of unnamed wood-decaying fungi have been reported previously (Orr and Charles, 1994). *Aphaenogaster* ants also visited and harvested pieces of basidiocarps of the genera

*Amanita*, *Armillariella*, *Marasmiellus*, and *Russula* (Carroll et al., 1981). Most recent findings showed laboratory and field evidences of interaction between wood-decaying fungi (*Laetiporus* and *Pleurotus ostreatus*) and ants *Aphaenogaster* species (Epps et al., 2018).

Decomposing dead pinewood provides essential habitat for various ants, beetles, and wood-decaying fungi (Warren and Bradford, 2011). Since many wood-decaying fungi are associated with various beetles and wasps, it raises the question whether they also have any sort of association with ants. It is challenging to prove the specific relationship of wood-decaying fungi and co-occurring ants because there can be various explanations for such co-occurrences: e.g., i) wood-decaying fungi may have a specific relationship with ants, ii) accidental co-occurrences, and iii) beetles may carry fungal spores and then ants use basidiocarps or mycelia as food source, etc. More research is needed to uncover such relationship.

## **5. Conclusions**

This study provides a new baseline in unravelling the complex interaction between wood-decaying fungi and ants in forest ecosystems. Both wood-decaying fungi and ants are important in providing nutrients to the soil through decomposition of wood debris; yet, much is not known on how they interact with each other at this specific niche of dead pinewood. Though this study is limited with small sample size, it nonetheless showed diverse patterns of co-occurrence between wood-decaying fungi and ants. Based on observations, future studies are needed to answer the following questions: Is there any specific interaction between wood-decaying fungi and ants? Do the ants participate in dispersal of their spores? If so, how do the mechanisms happened by functional way?

## 6. References

- Allen, R.B., Buchanan, P.K., Clinton, P.W., Cone, A.J., 2000. Composition and diversity of fungi on decaying logs in a New Zealand temperate beech (*Nothofagus*) forest. *Can. J. For. Res.* 30, 1025–1033.
- Breitenbach, J., Kränzlin, F., 1986. Fungi of Switzerland, Vol. 2. Non-gilled fungi. Verlag Mykologia, Lucerne.
- Carlile, M.J., Watkinson, S.C., Gooday, G.W., 2001. The Fungi (2nd ed.). Academic Press, San Diego, pp. 297–366.
- Carroll, J., Kimbrough, J., Whitcomb, W., 1981. Mycophagy by *Aphaenogaster* spp. (Hymenoptera: Formicidae). *Proc. Entomol. Soc. Wash.* 83, 326–331.
- Dai, Y.C., 2012. Polypore diversity in China with an annotated checklist of Chinese polypores. *Mycoscience.* 53, 49–80.
- Dong, M., 2017. The Ants of Korea. Nature and Ecology, Seoul.
- Ejrnæs, R., Bruun, H.H., Graae, B.J., 2006. Community assembly in experimental grassland, suitable environment or timely arrival? *Ecology.* 87, 1225–1233.
- Epps, M.J., Penick, C.A., 2018. Facultative mushroom feeding by common woodland ants (Formicidae, *Aphaenogaster* spp.). *Food Webs.* 14, 9–13
- Feldhaar, H., Fiala, B., Gadau, J., Mohamed, M., Maschwitz, U., 2003. Molecular phylogeny of *Crematogaster* subgenus *Decacrema* ants (Hymenoptera: Formicidae) and the colonization of *Macaranga* (Euphorbiaceae) trees. *Mol. Phylogenet. Evol.* 27(3), 441–452.
- Floren, A., Krüger, D., Müller, T., Dittrich, M., Rudloff, R., Hoppe, B., Linsenmair, K.E., 2015. Diversity and interactions of wood-inhabiting fungi and beetles after deadwood enrichment. *PLoS One.* 10(11), e0143566.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar. Biol. Biotechnol.* 3(5), 294–299.
- Fukasawa, Y., Matsuoka, S., 2015. Communities of wood-inhabiting fungi in dead pine logs along a geographical gradient in Japan. *Fungal Ecol.* 18, 75–82.

- Fukasawa, Y., Takahashi, K., Arikawa, T., Hattori, T., Maekawa, N., 2015. Fungal wood decomposer activities influence community structures of myxomycetes and bryophytes on coarse woody debris. *Fungal Ecol.* 14, 44–52.
- Gafforov, Y., Riebesehl, J., Ordynets, A., Langer, E., Yarasheva, M., Ghobad-Nejhad, M., Zhou, L.W., Wang, X.W., Gugliotta, A.M., 2017. *Hyphodontia* (Hymenochaetales, Basidiomycota) and similar taxa from Central Asia. *Botany*. 95, 1041–1056.
- Gardes, M., Bruns, T.D., 1994. ITS primers with enhanced specificity for basidiomycetes - application to the identification of mycorrhizae and rusts. *Mol. Ecol.* 2, 113–118.
- Hågvar, S., 1999. Saproxylic beetles visiting living sporocarps of *Fomitopsis pinicola* and *Fomes fomentarius*. *Nor. J. Entomol.* 46, 25–32.
- Heilmann-Clausen, J., Aude, E., Van Dort, K., Christensen, M., Piltaver, A., Veerkamp, M., Walley, R., Siller, I., Standovár, T., Òdor, P., 2014. Communities of wood-inhabiting bryophytes and fungi on dead beech logs in Europe – reflecting substrate quality or shaped by climate and forest conditions? *J. Biogeogr.* 41, 2269–2282.
- Herrmann, S., Bauhus, J., 2013. Effects of moisture, temperature and decomposition stage on respirational carbon loss from coarse woody debris (CWD) of important European tree species. *Scand. J. For. Res.* 28, 346–357.
- Hibbett, D.S., Binder, M., Bischoff, J.F., Blackwell, et al., 2007. A higher-level phylogenetic classification of the Fungi. *Mycol. Res.* 111(5), 509–547.
- Hinkle, G., Wetterer, J.K., Schultz, T.R., Sogin, M.L., 1994. Phylogeny of the Attini ants fungi based on analysis of small subunit ribosomal RNA gene sequences. *Science*. 266, 1695–1697.
- Hulcr, J., Stelinski, L.L., 2017. The ambrosia symbiosis: from evolutionary ecology to practical management. *Ann. Rev. Entom.* 62, 285–303.
- Jang, Y., Jang, S., Lee, J., Lee, H., Lee, H., Lee, Y.M., Hong, J.H., Min, M., Lim, Y.W., Kim, C., Kim, J.J., 2014. Wood decay fungi in South Korea: polypores from Seoul. *Mycobiology*. 42(2), 140–146.

- Jang, Y., Jang, S., Lee, J., Lee, H., Lim, Y.W., Kim, C., Kim, J.J., 2016. Diversity of wood-inhabiting polyporoid and corticioid fungi in Odaesan national park, Korea. *Mycobiology*. 44 (4), 217–236.
- Japanese Ant Image Database., 2010. <http://ant.edb.miyakyo-u.ac.jp/>.
- Johansson, T., Olsson, J., Hjältén, J., Jonsson, B.G., Ericson, L., 2006. Beetle attraction to sporocarps and wood infected with mycelia of decay fungi in old-growth spruce forests of northern Sweden. *For. Ecol. Manage.* 237(1), 335–341.
- Jonsell, M., Nordlander, G., 2004. Host selection patterns in insects breeding in bracket fungi. *Ecol. Entomol.* 29, 697–705.
- Jönsson, M., Edman, M., Jonsson, B.G., 2008. Colonization and extinction patterns of wood-decaying fungi in a boreal old-growth *Picea abies* forest. *J. Ecol.* 96, 1065–1075.
- Kim, C.S., Han, S.K., Nam, J.W., Jo, J.W., Kwag, Y.N., Han, J.G., Sung, G.H., Lim, Y.W., Oh, S., 2017. Fungal communities in a Korean red pine stand, Gwangneung forest, Korea. *J. Asia Pac. Biodiv.* 10(4), 559–572.
- Kim, G.H., Lim, Y.W., Choi, Y.S., Kim, M.J., Kim, J.J., 2009. Primary and secondary decay fungi on exposed pine tree logs in the forest. *Holzforschung*. 63, 633–638.
- Kim, G., Lyu, G., 2012. Distribution of ants (Insecta, Hymenoptera) in Chiaksan mountain, Prov. Gangweon, Korea. *J. Korean Nat.* 5(2), 127–129.
- Komonen, A., Ikävalko, J., Weiying, W., 2003. Diversity patterns of fungivorous insects: comparison between glaciated vs. refugial boreal forests. *J. Biogeogr.* 30, 1873–1881.
- Krasutskii, B.V., 2007. Beetles (Coleoptera) associated with the polypore *Daedaleopsis confragosa* (Bolton: Fr.) J. Schrot (Basidiomycetes, Aphyllophorales) in forests of the urals and transurals. *Entomol. Rev.* 87, 512–523.
- Kubartová, A., Ottosson, E., Dahlberg, A., Stenlid, J., 2012. Patterns of fungal communities among and within decaying logs, revealed by 454 sequencing. *Mol. Ecol.* 21, 4514–4532.

- Kula, E., Bohác, J., Jelínek, J., 1999. Insect fauna of selected polypore fungi on birch stems in northern Bohemia. *Misc. Zool*, 22, 75–85.
- Kwon, T.S., Kim, S.S., Lee, C.M., 2012. Korean ant atlas (2006-2009). Research Report 459. Korea Forest Research Institute. Seoul.
- Kwon, T.S., Park, Y.K., Lee, C.M., 2013. Change of ant communities in the burned forests in eastern coastal area. Research report 13-24. Korea Forest Research Institute. Seoul.
- Kwon, T.S., Kim S.S., Lee, C.M., 2014a. Pattern of ant diversity in Korea: An empirical test of Rapoport's altitudinal rule. *J. Asia Pac. Entomol.* 117, 161–167.
- Kwon, T.S., Lee, C.M., Sung, J.H., 2014b. Diversity decrease of ant (Formicidae, Hymenoptera) after a forest disturbance: different responses among functional guilds. *Zool. Stud.* 53, 37–47.
- Kwon, T.S., 2015a. Change of ant fauna in the Gwangneung long-term ecological research site in South Korea. *J. Asia Pac. Biodiv.* 8, 187–190.
- Kwon, T.S., 2015b. Foraging activity of competing ants along altitudinal gradient on a high mountain, South Korea. *J. Asia Pac. Entomol.* 19, 95–101.
- Kwon, T.S., Lee, C.M., 2015. Prediction of abundance of ants according to climate change scenarios RCP 4.5 and 8.5 in South Korea. *J. Asia Pac. Biodiv.* 8, 49–65.
- Kwon, T.S., 2016. What is important for ant assemblages in temperate forest soils? *J. Asia Pac. Biodiv.* 9, 138–143.
- Largent, D.L., Thiers, H.D., 1977. How to identify mushrooms to genus II: field identification of genera. Mad River Press, USA.
- Lee, J.S., Jung, H.S., 2006. Taxonomic study on Korean Aphyllophorales (5) - on some unrecorded genera and species. *Mycobiology*. 34(4), 166–175.
- Lewis, G.P., Worthen, W.B., 1992. Effects of ant predation and mushroom desiccation on the survival of mycophagous *Drosophila tripunctata* larvae. *Oikos*. 64, 553–559.

- Li, Y., Bateman, C.C., Skelton, J., Jusino, M.A., Nolen, Z.J., Simmons, D.R., Hulcr, J., 2017. Wood decay fungus *Flavodon ambrosius* (Basidiomycota: Polyporales) is widely farmed by two genera of ambrosia beetles. *Fungal Biol.* 121, 984–989.
- Li, Y., Simmons, D.R., Bateman, C.C., Short, D.P.G., Kasson, M.T., Rabaglia, R.J., Hulcr, J., 2015. New fungus-insect symbiosis: culturing, molecular, and histological methods determine saprophytic Polyporales mutualists of *Ambrosiodmus* ambrosia beetles. *PLoS One.* 10, e013768.
- Liu, W.D., Schaefer, L., Qiao, X.L., 2013. What controls the variability of wood-decay rates? *For. Ecol. Manag.* 310, 623–631.
- Liu, N., Yu, X., Xian, Z.Q., Wang, X.Y., Li, X.L., Ding, M.Y., 2016. Characterization of the complete mitochondrial genome of the myrmicine and *Vollenhovia emeryi* (Insecta: Hymenoptera: Formicidae). *Conservation Genet Resour.* 8 (3), 211–214.
- Lonsdale, D., Pautasso, M., Holdenrieder, O., 2008. Wood-decaying fungi in the forest: conservation needs and management options. *Eur. J. Forest Res.* 127(1), 1–22.
- Makonde, H.M., Boga, H.I., Osiemo, Z., Mwirichia, R., Stielo, J.B., 2013. Diversity of *Termitomyces* associated with fungus-farming Termites assessed by cultural and culture-independent methods. *PLoS One.* 8(2), e56464.
- Martin, K.J., Rygiewicz, P.T., 2005. Fungal-specific PCR primers developed for analysis of the ITS region of environmental DNA extracts. *BMC Microbiol.* 5, 28.
- Masiulionis, V.E., Rabeling, C., De Fine Licht, H.H., Schultz, T., Bacci, M.Jr., Cintia, M., Bezerra, S., Pagnocca, F.C., 2014. A Brazilian population of the asexual fungus-growing ant *Mycocepurus smithii* (Formicidae, Myrmicinae, Attini) cultivates fungal symbionts with gongylidia-like structures. *PLoS One.* 9(8), e103800.
- Mehdiabadi, N.J., Schultz, T.R., 2010. Natural history and phylogeny of the fungus-farming ants (Hymenoptera: Formicidae: Myrmicinae: Attini). *Myrmecol. News.* 13, 37–55.



- Mueller, U.G., Schultz, T.R., Currie, C.R., Adams, R.M., Malloch, D., 2001. The origin of the attine ant-fungus mutualism. *Q. Rev. Biol.* 76, 169–197.
- Mueller, U.G., Gerardo, N.M., Aanen, D.K., Six, D.L., Schultz, T.R., 2005. The evolution of agriculture in insects. *Annu. Rev. Ecol. Evol. Syst.* 36, 563–595.
- Nikitsky, N.B., Schigel, D.S., 2004. Beetles in polypores of the Moscow region: checklist and ecological notes. *Entomol. Fennica*. 15, 6–22.
- Olatinwo, R., Allison, J., Meeker, J., Johnson, W., Streett, D., Aime, M.C, Carlton, C., 2013. Detection and identification of *Amylostereum areolatum* (Russulales: Amylostereaceae) in the mycangia of *Sirex nigricornis* (Hymenoptera: Siricidae) in central Louisiana. *Environ. Entomol.* 42(6), 1246–1256.
- Orr, A., Charles, J., 1994 Foraging in the giant forest ant, *Camponotus gigas* (Smith) (Hymenoptera: Formicidae): evidence for temporal and spatial specialization in foraging activity. *J. Nat. Hist.* 28, 861–872.
- Osawa, N., Toft, R., Tuno, N., Kadowaki, K., Fukiharu, T., Buchanan, P.K., Tanaka, C., 2011. The community structures of fungivorous insects on *Amanita muscaria* in New Zealand. *NZ. Entomol.* 34, 40–44.
- Ottosson, E., Kubartová, A., Edman, M., Jonsson, M., Lindhe, A., Stenlid, J., Dahlberg, A., 2015. Diverse ecological roles within fungal communities in decomposing logs of *Picea abies*. *FEMS Microbiol. Ecol.* 91, 1–13.
- Park, M.S., Fong, J.J., Lee, H., Oh, S.Y., Jung, P.E., Min, Y.J., Seok, S.J., Lim, Y.W., 2013 Delimitation of *Russula* subgenus *Amoenula* in Korea using three molecular markers. *Mycobiology*. 41, 191–201.
- Park, M.S., Fong, J.J., Lee, H., Shin, S., Lee, S., Lee, N., Lim, Y.M., 2014. Determination of coleopteran insects associated with spore dispersal of *Cryptoporus volvatus* (Polyporaceae: Basidiomycota) in Korea. *J. Asia Pac. Entomol.* 17, 647–651.
- Pažoutová, S., Šrůtka, P., 2007. Symbiotic relationship between *Cerrena unicolor* and the horntail *Tremex fuscicornis* recorded in the Czech Republic. *Czech Mycol.* 59(1), 83–90.
- Persson, Y., Ihrmark, K., Stenlid, J., 2011. Do bark beetles facilitate the establishment of rot fungi in Norway spruce? *Fungal Ecol.* 4, 262–269.

- Rajala, T., Tuomivirta, T., Pennanen, T., Mäkipää, R., 2015. Habitat models of wood-inhabiting fungi along a decay gradient of Norway spruce logs. *Fungal Ecol.* 18, 48–55.
- Rogers, S.O., Bendich, A.J., 1994. Extraction of total cellular DNA from plants, algae and fungi. In: Gelvin S.B., Schilperoort R.A. (eds), *Plant molecular biology manual*. Kluwer Academic Publishers, Boston, pp 183–190.
- Schigel, D.S., 2011a. Fungus-beetle food web patterns in boreal forests. *Rus. Entomol. J.* 20, 141–150.
- Schigel, D.S., 2011b. Polypore–beetle associations in Finland. *Ann. Zool. Fennici.* 48, 319–348.
- Seibold, S., Müller, J., Baldrian, P., Cadotte, M.W., Štursová, M., Biedermann, P.H.W., Krah., F.Z., Bässler, C., 2019. Fungi associated with beetles dispersing from dead wood- Let's take the beetle bus! *Fungal Ecol.* 39, 100–108.
- Ševčík, J., 2003. Insects associated with wood-decaying fungi in the Czech and Slovak republics: a review of present knowledge. *Biol. Ecol.* 9, 159–165.
- Sippola, A.L., Mönkkönen, M., Renvall, P., 2005. Polypore diversity in the herb-rich woodland key habitats of Koli national park in eastern Finland. *Biol. Conserv.* 126, 260–269.
- Son, E., Kim, J.J., Lim, Y.W., Yeung, A., Yang, C.Y.H., Breuil, C., 2011. Diversity and decay ability of basidiomycetes isolated from lodge pole pines killed by the mountain pine beetle. *Can. J. Microbiol.* 57(1), 33–41.
- Strid, Y., Schroeder, M., Lindahl, B., Ihrmark, K., Stenlid, J., 2014. Bark beetles have a decisive impact on fungal communities in Norway spruce stem sections. *Fungal Ecol.* 7, 47–58.
- Takemoto, S., Nakamura, H., Imamura, E.Y., Shimane, K., 2010. *Schizophyllum commune* as a ubiquitous plant parasite. *JARQ.* 44 (4), 357–364.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.

- Tedersoo, L., Jairus, T., Horton, B.M., Abarenkov, K., Suvi, T., Saar, I., Kõljalg, U., 2008. Strong host preference of ectomycorrhizal fungi in a Tasmanian wet sclerophyll forest as revealed by DNA barcoding and taxon-specific primers. *New Phytol.* 180(2), 479–90.
- Thunes, K.H., Midtgaard, F., Gjerde, I., 2000. Diversity of coleopteran of the bracket fungus *Fomitopsis pinicola* in a Norwegian spruce forest. *Biodiversity Conserv.* 9, 833–852.
- Valer, F.B., Bernardi, E., Mendes, M.F., Blauth, M.L., Gottschalk, M.S., 2016. Diversity and associations between Drosophilidae (Diptera) species and Basidiomycetes in a Neotropical forest. *An. Acad. Bras. Ciênc.* 88, 705–718.
- Walker, J.K.M., Phillips, L.A., Jones, M.D., 2014. Ectomycorrhizal fungal hyphae communities vary more along a pH and nitrogen gradient than between decayed wood and mineral soil microsites. *Botany.* 92, 453–463.
- Warren, R.J., Bradford, M.A., 2011. Ant colonization and coarse woody debris decomposition in temperate forests. *Insect. Soc.* 59, 215.
- Weedon, J.T., Cornwell, W.K., Cornelissen, J.H., Zanne, A.E., Wirth, C., Coomes, D.A., 2009. Global meta-analysis of wood decomposition rates: a role for trait variation among tree species? *Ecol. Lett.* 12, 45–56.
- Weslien, J., Djupström, L.B., Schroeder, M., Widenfalk, O., 2011. Long-term priority effects among insects and fungi colonizing decaying wood. *J. Anim. Ecol.* 80, 1155–1162.
- White, T.J., Bruns, T., Lee, S.J., Taylor, J.W., 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications.* San Diego Academic Press, USA. pp 315–322.
- Wilson, J.J., 2012. DNA Barcodes for Insects. In: Kress W., Erickson D. (eds) *DNA Barcodes. Methods in Molecular Biology (Methods and Protocols)*, 858. Humana Press, New Jersey.
- Wooding, A.L., Wingfield, M.J., Hurley, B.P., Garnas, J.R., De Groot, P., Slippers, B., 2013. Lack of fidelity revealed in an insect-fungal mutualism after invasion. *Biol. Lett.* 9(4), 20130342.

- Wulandari, R., Lotrakul, P., Amirta, R., Kim, S.W., Punnapayak, H., Prasongsuk, H., 2018. First record of *Ceriporia inflata* and *Ceriporia lacerata* (Phanerochaetaceae, Basidiomycota) from Indonesian tropical forest. ANRES. 52, 412–418.
- Yamashita, S., Ando, K., Hoshina, H., Ito, N., Katayama, Y., Kawanabe, M., Maruyama, M., Itioka, K., 2015. Food web structure of the fungivorous insect community on bracket fungi in a Bornean tropical rain forest. Ecol. Entomol. 40, 390–400.
- Yu, E. J., Kim, T. H., Kim, K. H., Lee, H. J., 2004. Aroma-active compounds of *Pinus densiflora* (red pine) needles. Flavour Fragr. J 19, 532–537.

## 7. Abstract in Korean

### 한국의 죽은 소나무에서 공동 발생하는 목재부후균과 개미에 대한 고찰

루팔라

생명과학부 대학원

서울대학교

#### 초 록

죽은 소나무에 서식하는 개미, 딱정벌레, 말벌, 흰개미와 같은 곤충과 균류의 상호작용은 목재 잔해를 분해하고 분해 산물을 토양으로 환원할 수 있기 때문에 숲에서 생태학적으로 중요한 의미를 갖는다. 그러나 목재부후균과 개미 사이의 상호작용에 관한 정보는 매우 부족하다. 이 연구에서는 국내의 죽은 소나무에서 개미와 함께 발견되는 목재부후균을 조사하였으며, 각기 다른 여러 지역에서 총 57 쌍의 목재 부후균과 개미가 채집되었다. 형태 및 분자 동정에 근거하여 총 30 종의 목재부후균과 14 종의 개미가 확인되었다. *Trichaptum* 속, *Xylodon* 속, *Hyphodontia* 속, *Ceriporia* 속에 속한 균류가 우점하였으며, 여러 채집지에서 *Lasius* 속, *Camponotus* 속, *Pristomyrmex* 속,

*Crematogaster* 속에 속한 비교적 흔한 개미와 함께 채집되었다. 이 연구는 산림 생태계에서 목재부후균과 개미 사이의 복잡한 상호작용을 이해하기 위한 새로운 단초를 제시하였다 .

**주요어:** 개미, 공발생(共發生), 부식된 소나무, 적송

**학번:**2017-27643